

FIBER PRODUCTS OF HYPERELLIPTIC CURVES

SIMAN WONG

Let k be a number field, and let S be a finite set of maximal ideals of the ring of integers $\mathcal{O}_k$ of k . In his 1962 ICM address, Shafarevich [6] asked if there are only finitely many k -isomorphism classes of algebraic curves of a fixed genus $g \geq 1$ with good reduction outside S . He verified this for $g = 1$ by reducing the problem to Siegel's theorem. Parshin [5] extended this argument to all hyperelliptic curves (cf. also [4]). The general case was settled by Faltings' celebrated work [1]. In this note we give a short proof of Shafarevich's conjecture for hyperelliptic curves, by reducing the problem to the case $g = 1$ using the Theorem of de Franchis plus standard facts about discriminants of hyperelliptic equations.

Let $k, \mathcal{O}_k$ and S be as above. Enlarge S by a finite number of maximal ideals if necessary, so that

- (i) S contains every maximal ideal of $\mathcal{O}_k$ lying above 2; and
- (ii) the ring $\mathcal{O}_S$ of S -integers in k is a principal ring.

Denote by $k_{\mathfrak{p}}$ the residue field of a maximal ideal $\mathfrak{p}$ of $\mathcal{O}_S$. Denote by $\overline{F}$ a separable closure of a field F .

Fix an integer $g \geq 2$. Denote by $k(g, S)$ the compositum of all extensions of k of degree dividing $(2g + 2)!$ which are unramified outside S . By Hermite's theorem, $k(g, S)/k$ is a finite extension. To prove Shafarevich's Conjecture for hyperelliptic curves we proceed by induction on g , assuming as known the case $g = 1$. Following Lockhart [3], we say that a hyperelliptic curve over k is *pointed* if it has a k -rational Weierstrass point.

Denote by $\Sigma(k, g, S)$ the set of hyperelliptic curves over k of genus g with good reduction outside S . Define $\Sigma_0(k, g, S) = \{X \in \Sigma(k, g, S) : X \text{ is pointed over } k\}$.

Lemma. *Let $g \geq 2$. To prove that $\Sigma(k, g, S)$ is finite for every k and S , it suffices to prove that $\Sigma_0(k, g, S)$ is finite for every k and S .*

Proof. If X/k is a curve of genus ≥ 2 , then $\text{Aut}_{k(g, S)}(X)$ is finite, and hence so does $H^1(\text{Gal}(k(g, S)/k), \text{Aut}_{k(g, S)}(X))$. Thus there are at most finitely many k -isomorphism classes of curves over k which become isomorphic to X over $k(g, S)$. To prove the lemma it then suffices to show that the Weierstrass points of every curve $X \in \Sigma(k, g, S)$ are defined over $k(g, S)$.

Pick $X \in \Sigma(k, g, S)$. As 2 is a unit in $\mathcal{O}_S$ and X/k has good reduction outside S , for every maximal ideal $\mathfrak{p}$ of $\mathcal{O}_S$ the reduction mod $\mathfrak{p}$ map induces a bijection between the Weierstrass points of $X/\overline{k}$ and those of $X/\overline{k}_{\mathfrak{p}}$. Then the (conjugacy class of the) inertia group $I_{\mathfrak{p}}$ of $\mathfrak{p}$ in $\text{Gal}(\overline{k}/k)$ acts trivially on the Weierstrass points of $X/\overline{k}$, since $I_{\mathfrak{p}}$ acts trivially on $k_{\mathfrak{p}}$. There are exactly $2g + 2$ such points, so we are done. $\square$

Let $g \geq 2$. We now show that $\Sigma_0(k, g, S)$ is finite for all k and for all S satisfying (i) and (ii). Assume as inductive hypothesis that this is so for curves of positive genus $< g$, the case $g = 2$ being taken care of by Shafarevich [6]. Let X/k be a pointed hyperelliptic curve of genus g . By [3, Cor. 2.10] and by our choice of S , there exists a model for X of

the form $\mathcal{C}_X : y^2 + yQ(x) = P(x)$ with $Q, P \in \mathcal{O}_S[x]$, $\deg P = 2g + 1$, and $\deg Q \leq g$, such that for any maximal ideal $\mathfrak{p} \subset \mathcal{O}_k$ not in S , the curve X has bad reduction at $\mathfrak{p}$ if and only if the discriminant $\Delta(\mathcal{C}_X) := 2^{4g} \text{disc}(P(x) + \frac{1}{4}Q(x)^2)$ of the model $\mathcal{C}_X$ (cf. [3, §1]) is divisible by $\mathfrak{p}$, viewed as a maximal ideal in $\mathcal{O}_S$. Furthermore, since 2 is a unit in $\mathcal{O}_S$, by completing the square we can assume that $Q = 0$. So if X has good reduction outside S , then the principal ideal $\text{disc}(P)\mathcal{O}_S$ is equal to $\mathcal{O}_S$, and the polynomial $P \in k[x]$ splits completely over $K' = k(g, S)$. Without loss of generality we can assume that $P(0) \neq 0$. Let $R(z) = z^{2g+2}P(1/z)$. Then $y^2 = R(z)$ is another model of X over $\mathcal{O}_S$, we have $\deg R = 2g+2$, and R splits completely over K' . Thus we can write (not uniquely) $R = R_1 R_2$, where $R_i \in K'[x]$ and $\deg R_1 = 3, \deg R_2 = 2g - 1$. Denote by S' the set of primes of K' lying above those in S . From the discriminants of R_1 and R_2 we see that the two K' -rational hyperelliptic curves $X_i : y^2 = R_i(z)$ have good reduction outside S' . Moreover, both X_i are pointed over K' , and their genus are both positive and are less than that of C . By induction there are at most finitely many K' -isomorphism classes of such X_i/K' .

Denote by X_3/K' the fiber product $X_1 \times_{\mathbf{P}_{K'}^1} X_2$ with respect to the hyperelliptic involutions $X_i \rightarrow \mathbf{P}_{K'}^1$. An affine coordinate ring for X_3/K' is $A = K'[z, \sqrt{R_1}] \otimes_{K'[z]} K'[z, \sqrt{R_2}]$. It contains a subring isomorphic to $K'[z, \sqrt{R_1 R_2}]$, which is an affine coordinate ring for X/K' . Moreover, K' is algebraically closed in the field of fraction of A . Consequently, if we denote by X'_3/K' the smooth model of X_3/K' , there is a non-trivial K' -morphism $X'_3 \rightarrow X$. The Riemann-Hurwitz formula then implies that X'_3 has genus ≥ 2 since X does.

To recapitulate, with k, S and $g \geq 2$ as above, there exists a finite collection $\Xi(K', g, s)$ of smooth curves over K' of genus ≥ 2 , such that every $X \in \Sigma_0(k, g, S)$ when based changed to K' is the K' -image of some $X'_3 \in \Xi(K', g, S)$. By the Theorem of de Franchis [2], there are at most finitely many $\overline{K'}$ -isomorphism classes of pairs (π, Z) , where Z is a smooth projective curve of genus ≥ 2 over $\overline{K'}$, and $\pi : X'_3 \rightarrow Z$ is a non-trivial map over $\overline{K'}$. The same finiteness conclusion then holds if we replace $\overline{K'}$ by K' , since the automorphism group of any curve of genus ≥ 2 is finite. Consequently, $\Sigma_0(k, g, S)$ falls into finitely many K' -isomorphism classes. Invoke the H^1 -finiteness argument in the proof of the Lemma and we get the finiteness of $\Sigma_0(k, g, S)$ for all k and S . Apply the Lemma and we get the finiteness of $\Sigma(k, g, S)$ for all k and S , as desired.

REFERENCES

- [1] G. Faltings, Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. *Invent. Math.* **73** (1983) no. 3, 349–366.
- [2] M. de Franchis, Un teorema sulle involuzioni irrazionali. *Rend. Circ. Mat. Palermo* **36** (1913) 368.
- [3] P. Lockhart, On the discriminant of a hyperelliptic curve. *Trans. A. M. S.* **342** (1994) 729–752.
- [4] F. Oort, Hyperelliptic curves over number fields, in: *Classification of algebraic varieties and compact complex manifolds*, 211–218. LNM 412, Springer-Verlag, 1974.
- [5] A. N. Parshin, Minimal models of curves of genus 2, and homomorphisms of Abelian varieties defined over a field of finite characteristic. *Izv. Akad. Nauk SSSR Ser. Mat.* **36** (1972), 67–109. English trans: *Math. USSR: Izvestiya* **6** (1972) 65–109.
- [6] I. R. Shafarevich, Algebraic number fields, in: *Proc. Inter. Congress of Mathematicians*, 1962, 163–176. English trans: *A. M. S. Translations*. Ser. 2, vol. 31 (1963) 25–39.

DEPARTMENT OF MATHEMATICS & STATISTICS, UNIVERSITY OF MASSACHUSETTS. AMHERST, MA 01003-4515 USA

E-mail address: siman@math.umass.edu